ORIGINAL ARTICLE



WILEY Eve

Landscape genomics to the rescue of a tropical bee threatened by habitat loss and climate change

Rodolfo Jaffé^{1,2,3*} | Jamille C. Veiga^{4*} | Nathaniel S. Pope⁵ | Éder C. M. Lanes¹ | Carolina S. Carvalho¹ | Ronnie Alves¹ | Sónia C. S. Andrade⁶ | Maria C. Arias⁶ | Vanessa Bonatti⁷ | Airton T. Carvalho⁸ | Marina S. de Castro⁹ | Felipe A. L. Contrera⁴ | Tiago M. Francoy⁷ | Breno M. Freitas¹⁰ | Tereza C. Giannini¹ | Michael Hrncir³ | Celso F. Martins¹¹ | Guilherme Oliveira¹ | Antonio M. Saraiva¹² | Bruno A. Souza¹³ | Vera L. Imperatriz-Fonseca^{1,2,3}

⁴Instituto de Ciências Biológicas, Universidade Federal do Pará, Belém, Brazil

⁵Department of Integrative Biology, University of Texas, Austin, Texas

⁷Departamento de Genética, Faculdade de Medicina de Ribeirão Preto, Universidade de São Paulo, Ribeirão Preto, Brazil

⁸Unidade Acadêmica de Serra Talhada, Universidade Federal Rural de Pernambuco, Serra Talhada, Brazil

- ⁹Centro de Agroecologia Rio Seco, Universidade Estadual de Feira de Santana, Amélia Rodrigues, Brazil
- ¹⁰Departamento de Zootecnia, Universidade Federal do Ceará, Fortaleza, Brazil

¹¹Departamento de Sistemática e Ecologia, Universidade Federal da Paraíba, João Pessoa, Brazil

¹²Escola Politécnica da Universidade de São Paulo, Universidade de São Paulo, São Paulo, Brazil

¹³Embrapa Meio-Norte, Teresina, Brazil

Correspondence

Rodolfo Jaffé, Instituto Tecnológico Vale, Belém, Brazil. Email: r.jaffe@ib.usp.br

Funding information

Conselho Nacional de Desenvolvimento Científico e Tecnológico, Grant/Award Number: 300714/2017-3, 301616/2017-5, 302976/2015-9, 305126/2013-0, 306932/2016-4, 311531/2014-8 and 478982/2013-5; Research Center on Biodiversity and Computing; Empresa Brasileira de Pesquisa Agropecuária, Grant/ Award Number: 02.11.01.029.00.00; Instituto Tecnológico Vale; Coordenação de Aperfeiçoamento de Pessoal de Nível Superior, Grant/Award Number: 15/2014 and 88887.156652/2017-00; National

Abstract

Habitat degradation and climate change are currently threatening wild pollinators, compromising their ability to provide pollination services to wild and cultivated plants. Landscape genomics offers powerful tools to assess the influence of landscape modifications on genetic diversity and functional connectivity, and to identify adaptations to local environmental conditions that could facilitate future bee survival. Here, we assessed range-wide patterns of genetic structure, genetic diversity, gene flow, and local adaptation in the stingless bee *Melipona subnitida*, a tropical pollinator of key biological and economic importance inhabiting one of the driest and hottest regions of South America. Our results reveal four genetic clusters across the species' full distribution range. All populations were found to be under a mutation-drift equilibrium, and genetic diversity was not influenced by the amount of reminiscent natural

*These authors contributed equally.

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

 ${\ensuremath{\mathbb C}}$ 2019 The Authors. Evolutionary Applications published by John Wiley & Sons Ltd

¹Instituto Tecnológico Vale, Belém, Brazil

²Departamento de Ecologia, Universidade de São Paulo, São Paulo, Brazil

³Departamento de Biociências, Universidade Federal Rural do Semi-Árido, Mossoró, Brazil

⁶Departamento de Genética e Biologia Evolutiva, Universidade de São Paulo, São Paulo, Brazil

Science Foundation

volutionary Applications

WILEY

habitats. However, genetic relatedness was spatially autocorrelated and isolation by landscape resistance explained range-wide relatedness patterns better than isolation by geographic distance, contradicting earlier findings for stingless bees. Specifically, gene flow was enhanced by increased thermal stability, higher forest cover, lower elevations, and less corrugated terrains. Finally, we detected genomic signatures of adaptation to temperature, precipitation, and forest cover, spatially distributed in latitudinal and altitudinal patterns. Taken together, our findings shed important light on the life history of *M. subnitida* and highlight the role of regions with large thermal fluctuations, deforested areas, and mountain ranges as dispersal barriers. Conservation actions such as restricting long-distance colony transportation, preserving local adaptations, and improving the connectivity between highlands and lowlands are likely to assure future pollination services.

KEYWORDS

deforestation, environmental associations, gene flow, isolation by resistance, local adaptation, pollination, single nucleotide polymorphism, stingless bees

1 | INTRODUCTION

Although bees are now widely acknowledged as key pollinators of wild and cultivated plants, as well as important income sources for beekeepers around the globe (Potts et al., 2016), the joint impact of habitat degradation and climate change is currently threatening their wild populations (Brown & Paxton, 2009; Hadley & Betts, 2011; Potts et al., 2010; Viana et al., 2012; Wratten, Gillespie, Decourtye, Mader, & Desneux, 2012). Habitat loss has been related to reductions in native bee abundance and richness (Kennedy et al., 2013), whereas land use changes have fragmented populations in some species (Jha, 2015; Jha & Kremen, 2013). Additionally, climate change is expected to modify the availability of floral and nesting resources and affect bee physiology, thereby resulting in distribution range shifts and reductions in many species (Faleiro, Nemésio, & Loyola, 2018; Giannini et al., 2017; Kerr et al., 2015; Le Conte & Navajas, 2008; Pyke, Thomson, Inouye, & Miller, 2016; Willmer, 2014).

Landscape genomics offers powerful tools to assess the influence of habitat loss on genetic diversity and functional connectivity, and to identify adaptations to local environmental conditions that could facilitate future bee survival (Balkenhol et al., 2017; Lozier & Zayed, 2017). For instance, landscape resistance to gene flow has been assessed in both temperate and tropical species (Davis, Murray, Fitzpatrick, Brown, & Paxton, 2010; Jackson et al., 2018; Jaffé, Castilla, et al., 2016), and genomic signatures of adaptations to environmental conditions have been identified in the honeybee *Apis mellifera* (Chávez-Galarza et al., 2013; Henriques et al., 2018) and the bumblebee *Bombus lapidarius* (Theodorou et al., 2018). Nonetheless, while most studies assessing landscape effects on gene flow have employed microsatellite markers (Balkenhol, Cushman, Waits, & Storfer, 2016; Monteiro et al., 2019), none has thus far employed genomic data to assess both isolation by landscape resistance and local adaptation in bees (Storfer, Patton, & Fraik, 2018).

Dispersal is believed to be particularly restricted in stingless bees (Apidae: Meliponini), because daughter colonies rely on resources from their maternal colonies during their initial establishment and consequently do not establish far from each other (Roubik, 2006; Van Veen & Sommeijer, 2000; Vit, Pedro, & Roubik, 2013). As restricted dispersal implies a diminished ability to move to high-quality habitats and maintain gene flow across fragmented landscapes, habitat loss and fragmentation are expected to reduce and isolate stingless bee populations, making them extremely susceptible to genetic erosion through the action of genetic drift (Allendorf, Luikart, & Aitken, 2013; Lozier & Zayed, 2017). However, previous studies using microsatellite markers were not able to detect an effect of forest or land cover on stingless bee gene flow, suggesting that these bees have a remarkable ability to maintain high gene flow across heterogeneous and human-altered landscapes (Jaffé, Castilla, et al., 2016; Jaffé, Pope, et al., 2016; Landaverde-González et al., 2017). Since estimates of genetic diversity and gene flow are strongly influenced by the type and number of genetic markers employed (Allendorf, 2017; Leroy et al., 2018; Lozier, 2014), genomic studies employing thousands of single nucleotide polymorphisms (SNPs) are needed to confirm whether stingless bees are really resilient to habitat loss and fragmentation, or whether the lack of significant isolation-by-resistance effects in previous studies is due to the resolution of the genetic markers employed (Alvarado-Serrano, Van Etten, Chang, & Baucom, 2019; McCartney-Melstad, Vu, & Shaffer, 2018).

Here, we employ novel landscape genomic tools to assess the joint influence of habitat degradation and climate change on an economically important tropical stingless bee inhabiting one of the most deforested, driest, and hottest regions of the Americas. Distributed across northeastern Brazil, *Melipona subnitida* Ducke, 1911 (known WILEY

Evolutionary Applications

as the Jandaíra bee) is a key pollinator of native plants and local crops and one of the most widely used stingless bee species for honey production (Imperatriz-Fonseca, Koedam, & Hrncir, 2017), contributing to the household income of many rural families (Giannini et al., 2017; Jaffé et al., 2015). The species natural range spans four different biomes (Giannini et al., 2017; Imperatriz-Fonseca et al., 2017), namely Tropical Dry Forest (Caatinga), Savanna (Cerrado), the Atlantic Rain Forest, and Mangrove Forests, thus encompassing important climatic and altitudinal gradients, which are expected to drive local adaptations (Koffler et al., 2015; Maia-Silva, Hrncir, Silva, & Imperatriz-Fonseca, 2015). Species distribution models suggest the species could respond to climate change by seeking refuge in higher elevations, where both the bees and their plant resources are more likely to find suitable climatic conditions in the future (Giannini et al., 2017). However, such response depends on the bee's ability to relocate to high-quality habitats, which could be hindered by the region's increasing human-led desertification (Marengo, Torres, & Alves, 2017; Vieira et al., 2015). Given M. subnitida's key biological and economic importance, efforts are urgently needed to safeguard this key pollinator by facilitating its migration toward higher lands. Such conservation actions will nevertheless require the prior identification of barriers to gene flow and the spatial distribution of adaptive genetic variation.

Relying on thousands of SNPs, we explicitly tested whether genetic diversity and gene flow in *M. subnitida* are affected by the amount of reminiscent natural habitats, and identified genomic signatures of adaptations to local environmental conditions. Considering the life history characteristics of our study species and the high statistical power granted by the large number of genetic markers employed, we formulated the following predictions: (a) Since natural habitats harbor floral resources and nesting sites (Vit et al., 2013), we expected to find a positive association between genetic diversity and the amount of reminiscent natural habitats surrounding sampling sites (DiLeo & Wagner, 2016;

Fahrig, 2013), as found in bumblebees (Jackson et al., 2018); (b) as restricted dispersal implies reduced gene flow across deforested areas, we expected to find significant isolation by landscape resistance (McRae, 2006) and predicted that gene flow would be influenced by habitat amount as well as environmental correlates of genetic connectivity in other bee species, including elevation. terrain roughness, temperature, and precipitation (El-Niweiri & Moritz, 2011; Jackson et al., 2018; Jaffé, Pope, et al., 2016; Jha, 2015); and (c) based on the documented tolerance of our study species to extreme heat and water scarcity (Maia-Silva et al., 2015), and previous candidate genes found associated with precipitation and latitude in honeybees (Chávez-Galarza et al., 2013; Henriques et al., 2018) and urban land cover in bumblebees (Theodorou et al., 2018), we expected to find genomic signatures of adaptation related to temperature, precipitation, and forest cover. Ours constitutes the first genomic study assessing both isolation by landscape resistance and local adaptation across the full distribution range of a bee pollinator.

2 | MATERIALS AND METHODS

2.1 | Sampling and DNA extraction

We collected samples of *M. subnitida* across its entire distribution range (Pedro, 2014), aiming to maximize temperature, precipitation, elevation, and forest cover gradients (Figure 1). Samples were collected between 2013 and 2014 (SISBIO collection permits 38000-1 and 10393-1). We sampled one bee per colony from a total of 160 nests of beekeepers who could certify their local origin. We only collected samples from local colonies, and not from colonies of unknown origin or from nests brought from different locations. We also registered any information on previous introductions of bee colonies that beekeepers could provide, as long-distance colony transportation is common and has been shown to have profound effects on



FIGURE 1 Melipona subnitida sampling locations across northeastern Brazil over a land cover map (source: http:// mapbiomas.org/)

-WILEY

stingless bee gene flow (Jaffé, Pope, et al., 2016). In cases when beekeepers had a few weak colonies, we did not collect any samples or collected samples from a single colony. When beekeepers had many strong colonies, we collected samples from more than one colony. Distance separating sampling locations therefore ranged between 0 km (samples from the same beekeeper) and 947 km. Only freshly emerged (callow) workers were collected from the nest's interior, to minimize the sampling of drifters. All individuals were stored in absolute ethanol and then frozen at -20° C until DNA extractions. The geographic location of all samples was recorded by GPS (see Supporting Information Table S1).

Total genomic DNA was extracted from the whole insects (discarding only the heads) using Qiagen's DNeasy Blood and Tissue Kit, according to the manufacturer's protocols. DNA integrity was then verified on 1% agarose gels, and its concentration was quantified with a Qubit 2.0 Fluorometer (Invitrogen). Only samples containing nondegraded DNA, concentrations of at least 20 ng/µl, and a total DNA amount greater than 2 µg were used for subsequent RAD sequencing.

2.2 | RAD sequencing and SNP discovery

DNA samples were then shipped to Floragenex, Inc. (Eugene, OR, USA) for RAD library preparation, Illumina sequencing, and bioinformatic processing. Briefly, libraries were prepared based on the genome size of this species (Tavares, Carvalho, Aparecida, & Soares, 2010), digesting genomic DNA with the SgrAl restriction enzyme. The resulting fragments were tagged with individual barcodes, which were then multiplexed and sequenced using 100-bp singleend methodology on the HiSeq 2000 platform (Illumina). Libraries were created at the same time and ran on the same sequencing run on two different lanes. Total number of generated reads per individual ranged from ~550,000 to 2.3 million. Samples were then demultiplexed and barcode sequences trimmed to result in final fragment lengths of 92 bp. Quality filtering was done in SAMtools during the genotyping stages, and coverage was limited to loci having less than 500× to limit the incidence of possible contaminants. Stacks was then used to cluster loci from a single individual and generate a RAD reference, allowing two haplotypes from each locus. The de novo clustering of sequences into RAD tags was performed using VELVET (version 1.2.10), considering a minimum cluster depth of 5 and maximum of 1,500, a maximum number of two haplotypes per cluster, and a maximum of three variants per cluster. SNP calling was performed using SAMtools (version 0.1.16), and loci harboring SNPs were included in the final genotype (VCF) table if they had at least 6× individual sequence coverage over at least 75% of the population, individual per locus genotype quality scores of at least 10, a minimum FASTQ quality score of 20, and a minimum distance to other SNPs of 50 bp (average individual Phred score was 60.8, while average individual sequencing coverage was 36.3×). A variant was cataloged when it was present in a single sample (only 3% of loci had minor allele frequencies-MAF-below 0.5, 2% of loci had MAF below 0.3, and 1% of loci had MAF below 0.1.).

2.3 | Population structure and genetic diversity

The R package *r2vcftools* (https://github.com/nspope/r2vcftools), a wrapper for VCFtools (Danecek et al., 2011), was used to perform final filtering and quality control on the genotype data. To assess genetic diversity and population structure across the distribution range of our study species, we first filtered loci for quality (Phred score 30–80), read depth (20–50), linkage disequilibrium (LD, $r^2 < 0.4$), and strong deviations from the Hardy–Weinberg equilibrium (HWE, p < 0.0001). Additionally, we removed any potential loci under selection detected through genome scans. F_{ST} outlier tests were applied after adjusting *p*-values using the genomic inflation factor (λ), and setting false discovery rates to q = 0.05, using the Benjamini–Hochberg algorithm (Benjamini & Hochberg, 1995; François, Martins, Caye, & Schoville, 2016).

Two complementary genetic clustering software packages were employed to assess population structure using the resulting set of neutral and independent loci: the *snmf* function of the LEA (v2.0) package (Frichot & François, 2015; Frichot, Mathieu, Trouillon, Bouchard, & François, 2014) and Admixture (Alexander, Novembre, & Lange, 2009). The number ancestral populations (k) was allowed to vary between 1 and 10, with 10 replicate runs for each k-value, and the best k was chosen based on cross-entropy and cross-validation errors (Frichot et al., 2014). Individuals were then assigned to genetic clusters based on the ancestry coefficients retrieved from LEA (Qmatrix), identifying the cluster with the highest ancestry.

We then calculated genetic diversity metrics for each one of the identified genetic clusters. These included observed heterozygosity (H_{Ω}) , expected heterozygosity (H_{F}) , nucleotide diversity (π) , and inbreeding coefficient (F; see VCFtools manual for details). Effective population size (N_{a}) was also estimated employing the linkage disequilibrium method implemented in NeEstimator 2.0.1 (Do et al., 2014), using a threshold lowest allele frequency value of 0.05 and assuming a monogamy model (Jaffé et al., 2014). Additionally, we calculated Tajima's D, representing the difference between the mean number of pairwise differences and the number of segregating sites. In a population of constant size evolving under mutation-drift equilibrium, Tajima's D is expected to be zero; negative values result from an excess of rare alleles and thus indicate a recent selective sweep or a population expansion after a recent bottleneck; and positive values appear when rare alleles are lacking, therefore suggesting balancing selection or a sudden population contraction (Tajima, 1989). We used r2vcftools to compute a genome-wide estimate of Tajima's D and perform a simulation from the neutral model to correct for bias due to a minor-allele-frequency filter.

2.4 | Landscape genetic analyses

Aiming to assess the influence of habitat amount on genetic diversity, we reclassified a high-resolution land cover/land use map for 2013 (http://mapbiomas.org/) into habitat (i.e., all types of natural forest and nonforest formations) and nonhabitat (farming, nonvegetated areas, and water bodies). The percentage of habitat cover in NILEY-

Evolutionary Applications

2-km-radius buffers surrounding our sampling locations was then calculated, as such radius comprises the estimated foraging distance for this species (Silva & Ramalho, 2016). Because more than one colony was sampled in some locations (Supporting Information Table S1), we computed mean percentage of habitat cover and mean genetic diversity (H_{Ω} , H_{F} , and F) for each location (N = 56 locations). We then used the nlme package (Pinheiro, Bates, DebRoy, & Sarkar, 2018) to fit generalized least squares models (gls) containing genetic diversity metrics as response variables, percentage of habitat cover as predictor, and different correlation structures (no autocorrelation, linear, exponential, Gaussian, spherical, and rational quadratics) to account for spatial autocorrelation. Logit transformations were used to normalize/linearize heterozygosities. The sample-size-corrected Akaike information criterion (AICc) was then used to compare models with different correlation structures, fitted with restricted maximum likelihood. The best models ($\Delta AICc \leq 2$) were finally selected, fitted once more using maximum likelihood, and compared to reduced models without predictor variables using likelihood ratio tests (LRT, α = 0.05). All models were validated by plotting residual versus fitted values and by checking for residual autocorrelation.

Prior to assessing isolation by landscape resistance (IBR), we evaluated fine-scale spatial genetic structure by quantifying spatial autocorrelation in genetic relatedness. To do so, we used local polynomial fitting (LOESS) of pairwise relatedness to pairwise geographic distance (https://github.com/rojaff/Lplot; Bruno, Macchiavelli, & Balzarini, 2008). Yang's relatedness between pairs of individuals (Yang et al., 2010) was used, since similar measures of relatedness have been found to be highly accurate as individual-based genetic distance metrics for landscape genetic studies (Shirk, Landguth, & Cushman, 2017). We then evaluated the contribution of habitat amount, elevation, terrain roughness, temperature, and precipitation in explaining patterns of gene flow (Balkenhol et al., 2016). We created a first resistance surface using the reclassified land cover/ land use map for 2013 described above, attributing low resistance (0.1) to habitat pixels and high resistance (0.9) to nonhabitat pixels. This allowed us to test whether gene flow is enhanced by natural habitats or hindered by habitat-deployed environments (Jaffé, Castilla, et al., 2016). A second resistance surface was then created using an inverted forest cover map from the University of Maryland (http://earthenginepartners.appspot.com/science-2013-global-forest/download.html), to test for higher gene flow across forested areas. To test for a reduced gene flow across highlands and corrugated terrains, we created elevation and terrain roughness surfaces, using raw elevation and terrain roughness maps. While elevation was retrieved from WorldClim (http://www.worldclim.org/), terrain roughness was created from this elevation layer using the Terrain Analysis plug-in in QGIS V2.14. To evaluate resistance due to environmental conditions, we created three additional resistance surfaces containing the raw values from those WorldClim bioclimatic variables explaining most variation across our study region (mean temperature of coldest quarter, temperature annual range, and precipitation of driest quarter; see details below). We thereby tested for reduced gene flow across areas with higher temperatures, higher

temperature range, and higher precipitation. To assess isolation by geographic distance (IBD), we created a last resistance surface replacing all pixel values in our elevation map with 0.5. Using the program Circuitscape V4.0 (McRae, 2006), we then calculated pairwise resistance distances between all samples, employing all the resistance surfaces described above. Due to Circuitscape's computing limitations, all rasters were cropped to the extent of sample locations plus a buffer area of one decimal degree to minimize border effects (Jaffé, Castilla, et al., 2016), and all pixels containing zero values were replaced with 0.001.

To assess IBR, we fit mixed-effects regression models using penalized least squares and a novel correlation structure designed to account for the nonindependence of pairwise distances across individuals and spatial locations (based on the maximum-likelihood population effects or MLPE model: https://github.com/nspope/ corMLPE; Clarke, Rothery, & Raybould, 2002). Because more than one colony was sampled in some locations (Supporting Information Table S1), there exists a spatial dependence structure in pairwise comparisons between individuals that is not captured by the MLPE model. This is because the MLPE correlation structure only models dependence across pairwise comparisons that overlap in the individuals being compared, with no reference to spatial location. The unmodeled spatial dependence has the net effect of making inference anticonservative and can be diagnosed by calculating serial autocorrelation in the normalized residuals of the MLPE model, after sorting the data by spatial locations. To combat this, we introduce a novel modification of the MLPE model that incorporates correlation between pairwise measurements due to comparison of both individuals and spatial locations (Nested MLPE or NMLPE). Briefly, the original MLPE model has a random-effects representation where the expected value for each pairwise observation includes a pair of iid random effects (one for each of the individuals being compared in the observation), whereas our extension (NMLPE) additionally incorporates iid random effects for pairs of spatial locations (see Supporting Information Script S1 and Data S1).

Yang's relatedness between pairs of individuals was used as response variable and the different resistance distances (geographic distance, forest cover, elevation, roughness, temperature annual range, mean temperature of coldest quarter, and precipitation of driest quarter) as predictors in our MLPE models. We used the Akaike information criterion (AIC) to compare models containing all possible combinations of noncollinear predictors (r < 0.6), created with the dredge function from the MuMIn (v1.4) package (https://github.com/ rojaff/dredge_mc; Barton, 2018). Likelihood ratio tests were then performed to assess the influence of the inclusion of each predictor variable on the best-fitting model's log-likelihood. Finally, we refitted the best models using NMLPE models to obtain parameter estimates unbiased by spatial dependence. To evaluate the impact of excluding samples from our IBR analyses, we also ran a sensitivity analysis, generating one hundred data subsets (randomly excluding different numbers of samples) and performing one hundred independent model selection protocols for each subset (using MLPE models). We report the number of times predictor variables were included

in the set of best-fitting MLPE regression models ($\Delta AIC \le 2$), after randomly excluding different numbers of samples (Supporting Information Figure S5).

2.5 | Identification of putative adaptive loci

To identify genomic signatures of adaptations to local environmental conditions, we employed environmental association tests. To this end, we used our original dataset filtered by quality and depth only (as described above), but not for LD or HWE. We first ran a principal component analysis using all 19 WorldClim bioclimatic variables plus altitude and forest cover (all scaled), to select a set of orthogonal variables explaining most environmental variation across our study area. Since the first four principal components accounted for 92.08% of total variance, we selected the four variables that were most strongly correlated with these axes. The selected variables (mean temperature of coldest guarter, forest cover, temperature annual range, and precipitation of driest quarter) were then used to run latent factor mixed models (LFMMs), aiming to identify possible associations between SNPs and environmental variables, while accounting for the underlying population structure (De Kort et al., 2014; Frichot, Schoville, Bouchard, & François, 2013; Rellstab, Gugerli, Eckert, Hancock, & Holderegger, 2015). LFMMs have been used extensively and are currently one of the most commonly used environmental association analysis approaches (Ahrens et al., 2018), given they provide a good compromise between detection power and error rates, and are robust to a variety of sampling designs and underlying demographic models (Rellstab et al., 2015). LFMM were implemented in R through the LEA package, using 1,000 iterations, a burn-in of 5,000, and five runs per environmental variable (Frichot & François, 2015). The *p*-values were adjusted using the genomic inflation factor (λ = 1), and false discovery rates were set using the

Benjamini–Hochberg algorithm at a rate of q = 0.05 (Benjamini & Hochberg, 1995). Since incorrect assumptions about underlying demographic structure can increase both type I and type II errors (Cushman & Landguth, 2010; Storfer et al., 2018), we ran LFMM using $k \pm 1$ latent factors (where k was the optimum number of ancestral populations detected) and only considered as candidate loci those shared between all runs for each environmental variable. Full R scripts of LFMM can be found in the LEA website (http://membres-timc.imag.fr/Olivier.Francois/LEA/index.htm). In order to map adaptive genetic variability, we used the *adegenet* package (Jombart, 2008) to run a spatial principal component analysis (sPCA) on all identified candidate SNPs and interpolated the first two principal components on a grid covering our study area.

3 | RESULTS

3.1 | Genetic diversity and population structure

We identified 29,349 SNPs from which 3,454 loci remained after filtering for quality, depth, HWE, LD, and F_{ST} outlier loci. We detected four genetic clusters (K = 4) using two different clustering approaches (Figure 2, Supporting Information Figure S1), and excluded four samples from subsequent analyses (final sample size was 156 individuals) because they were likely introduced bees from a distant location (Supporting Information Figure S2, Table S1). Cluster assignments were generally unambiguous as ancestry coefficients for the assigned clusters were usually above 0.50 (lower 25% quantile = 0.50, median = 0.60, upper 75% quantile = 0.86). Clusters 1 and 4, located at the northern and southern extremes of the species distribution range, showed the lowest N_e and π . All genetic clusters showed small but significant inbreeding and values of Tajima's *D* overlapping zero (Table 1).



FIGURE 2 Map showing *Melipona* subnitida assignments to four genetic clusters against an elevation map (from USGS Earth Explorer). Pie charts represent ancestry coefficients determined using the LEA package

ILEY Evolutionary Applications

TABLE 1 Genetic diversity estimates for Melipona subnitida by genetic cluster

Notes. Sample sizes (N) are shown followed effective population size (N_e), observed heterozygosity (H_o), expected heterozygosity (H_E), inbreeding coefficient (F), nucleotide diversity (π), and Tajima's D. Lower and upper 95% confidence intervals are shown for each estimate.

3.2 | Landscape genetic analyses

Habitat amount was not found associated with heterozygosity or inbreeding (Table 2). We found positive spatial autocorrelation in pairwise genetic relatedness for up to 300 km, after which spatial autocorrelation became negative (Figure 3). However, IBR was better able to explain genetic relatedness patterns than IBD (Table 3). Temperature annual range (defined as the difference between maximum temperature of warmest month and minimum temperature of coldest month) was the best predictor of relatedness patterns across the full distribution range of M. subnitida. The second best IBR model was nearly 18 AIC units apart from the model containing temperature annual range (Table 3), suggesting that forest cover, altitude, and terrain roughness combined did not explain relatedness patterns as well as temperature fluctuations alone. All these predictors showed a significantly negative association with genetic relatedness (Table 4, Figure 4). Although temperature annual range did not show a larger variation than the other predictors used to assess IBR (Supporting Information Figure S3), it was highly correlated with forest cover (r = 0.94; Supporting Information Figure S4), so the effect of temperature annual range is likely confounded by forest cover to some extent. Nested MLPE models (accounting for spatial dependence) substantially improved fit and reduced evident autocorrelation. Although parameter estimates from NMLPE models had slightly larger standard errors (implying these were more conservative estimates), all effects continued to be significantly different from zero (Table 4). Our sensitivity analysis on IBR models revealed that temperature annual range, altitude, and terrain roughness were the most frequent variables included in the set of best models (Supporting Information Figure S5).

3.3 | Identification of putative adaptive loci

About 10% of all analyzed sequences contained signatures of selection (Table 5), and most of the identified candidate loci were associated with temperature (Figure 5). The spatial distribution of adaptive genetic variability revealed latitudinal and altitudinal gradients (Figure 6).

4 | DISCUSSION

Our study reveals a clinal change in genetic structure across the distribution range of *M. subnitida*, with four identifiable genetic clusters.

TABLE 2	Effect of habitat amo	unt on observed	heterozygosity
$(H_{\rm O})$, expected	ed heterozygosity (H _E)	, and inbreeding	coefficient (F)

Response variable	Correlation structure	X ²	p-Value
H _o	Exponential	0.24	0.63
H _E	None	0.22	0.64
F	Exponential	0.25	0.62

Notes. Generalized least squares models contained genetic diversity metrics as response variables, percentage of habitat cover as predictor, and different correlation structures to account for spatial autocorrelation. Logit transformations were used to normalize/linearize heterozygosities. The table shows X^2 values and *p*-values from likelihood ratio tests applied on best-fitting models.



FIGURE 3 Spatial autocorrelation in genetic relatedness. The black solid line is the LOESS fit to the observed genetic relatedness, while the gray shaded regions are 95% confidence bounds around the null expectation (black dotted line). Short vertical lines at the bottom of the figure are observed pairwise distances

Genetic diversity was not influenced by habitat amount, pairwise relatedness showed spatial autocorrelation, and isolation by resistance explained range-wide relatedness patterns better than isolation by geographic distance. Specifically, gene flow was enhanced by low annual temperature variation, more forest cover, lower elevations, and flatter terrains. Finally, we detected genomic signatures of

TABLE 3 Summary statistics for the top MLPE regression models

Predictors	logLik	AIC	ΔAIC	Weight	ρ
Temperature annual range ^{***}	22,813.91	-45,619.8	0.00	1	0.27
Inverted forest cover***, altitude***, terrain roughness*	22,807.03	-45,602.1	17.76	0	0.23
Inverted forest cover***	22,576.63	-45,145.3	474.56	0	0.25
Geographic distance	22,521.73	-45,035.5	584.37	0	0.27

1171

Notes. All models contained interindividual genetic relatedness as response variable and the different landscape resistance distances as predictors. Log-likelihoods are followed by the Akaike information criterion (AIC), Δ AIC, model weight, and the MLPE correlation coefficient rho (ρ). Isolation by geographic distance was included here for comparison. Likelihood ratio tests: *p < 0.05, **p < 0.01, *p < 0.001

TABLE 4Parameter estimates for the
best-fitting MLPE regression models
 $(\Delta AIC < 20; see Table 3)$ and NMLPE
regression models (unbiased by spatial
dependence; in parentheses)

Predictors	Estimate	SE	CI
Temperature annual range	-0.14 (-0.09)	0.001 (0.003)	-0.14/-0.14 (-0.1/-0.09)
Inverted forest cover	-0.11 (-0.07)	0.002 (0.005)	-0.12/-0.11 (-0.08/-0.06)
Altitude	-0.02 (-0.01)	0.001 (0.003)	-0.02/-0.02 (-0.01/-0.001)
Terrain roughness	-0.01 (-0.02)	0.002 (0.005)	-0.01/-0.001 (-0.03/-0.01)
Geographic distance	-0.14 (-0.15)	0.001 (0.001)	-0.14/-0.14 (-0.1/-0.08)

Notes. Estimates are followed by standard errors (*SE*) and 95% confidence intervals (CI). Although the isolation by geographic distance was not among the top models, we include it here for comparison.



FIGURE 4 Isolation-by-resistance effects across the entire distribution range of *Melipona subnitida*. Plots show the relationship between genetic relatedness and temperature annual range (a), inverted forest cover (b), altitude (c), terrain roughness (d), and geographic distance (e). Although the isolation by geographic distance was not among the top models, we include it here for comparison. Relatedness is decorrelated for the MLPE correlation structure

adaptation to temperature, precipitation, and forest cover and found latitudinal and altitudinal patterns in the spatial distribution of adaptive genetic variation. Although previous genetic studies found intraspecific variation and signals of population structure in *M. subnitida* (Bonatti, Simões, Franco, & Francoy, 2014; Cruz et al., 2006; Silva et al., 2014), we Evolutionary Applications

TABLE 5 Summary of the number of adaptive signals detected employing environmental association tests. Both the number of candidate SNPs and the number of contigs (RAD tags) containing candidate SNPs are presented for each environmental predictor followed by the number of independent (nonoverlapping) detections in parentheses

		Total under	Environmental association tests ^a			
Signal type Total analyzed	Total analyzed	selection	Mean Temp CoQ	Forest Cover	Temp AnR	Prec DrQ
SNPs	27,799	1,798	997 (444)	718 (281)	700 (261)	478 (67)
Contigs	15,924	1,356	768 (334)	532 (195)	535 (203)	371 (45)

^aEnvironmental variables: mean temperature of coldest quarter (Mean Temp CoQ), forest cover, temperature annual range (Temp AnR), and precipitation of driest quarter (Prec DrQ).

here present the first assessment of spatial genetic structure based on thousands of independent genetic markers and performed across the full distribution range of this species. Our results reveal four genetic clusters, confirmed by two complementary methods, and extensive admixture (Figure 2). Genetic clusters located at the northernmost and southernmost extremes of the species' distribution range (Pop 1 and Pop 4) showed lower nucleotide diversity and lower effective population size (N_e) but similar heterozygosity and inbreeding as the genetic clusters located at the distribution's core (Pop 2 and Pop 3). This finding suggests that *M. subnitida* colonized these peripheral regions more recently, whereas clusters Pop 2 and Pop 3 had more time to accumulate a higher genetic variability, as stated by the central-peripheral hypothesis (Diniz-Filho et al., 2009).

Genetic diversity was slightly higher than that reported in other genomic bee studies (Jackson et al., 2018; Romiguier et al., 2014), and in all genetic clusters we found Tajima's D values overlapping with zero, which indicate mutation-drift equilibrium. Additionally, we did not find a significant association between habitat amount and genetic diversity. These results suggest that genetic variation has not been influenced by habitat loss yet and that the observed levels of inbreeding are presumably related to the reproductive biology of this species (DiLeo & Wagner, 2016). Monogamy (Jaffé et al., 2014), low population densities, and sharp seasonal variations in the production of reproductive individuals (Ferreira, Blochtein, & Serrão, 2013; Roubik, 2006; Santos-Filho, Alves, Eterovic, Imperatriz-fonseca, & Kleinert, 2006) may result in mating between related individuals. We nevertheless caution that longer time lags may be necessary to detect an effect of habitat loss on genetic diversity (Schlaepfer, Braschler, Rusterholz, & Baur, 2018).

We detected significant spatial autocorrelation in genetic relatedness, contradicting earlier microsatellite-based studies for other stingless bee species (Jaffé, Castilla, et al., 2016; Landaverde-González et al., 2017). Interestingly, spatial autocorrelation was positive for up to 300 km (Figure 3), indicating that related colonies can be found across large areas and that gene flow did not erase the genetic signals left by limited colony dispersal. Above 300 km, spatial autocorrelation became negative, suggesting that population differentiation caused lower than random relatedness between individuals from different clusters (Figure 2). These results suggest it would be safe to transport colonies no further than 300 km to avoid altering the genetic composition of wild populations (Jaffé, Pope, et al., 2016).

Isolation by landscape resistance explained range-wide relatedness patterns better than isolation by geographic distance alone. For instance, temperature fluctuations were found to be the most important factor explaining relatedness patterns in M. subnitida, followed by forest cover, elevation, and terrain roughness. Our IBR results hold when accounting for spatial dependence and when excluding different numbers of samples, although forest cover looses importance when more samples are excluded (Supporting Information Figure S5). Significant isolation by resistance (altitude) was only found in one other stingless bee (Partamona helleri Friese, 1900), out of 18 analyzed species to date (Jaffé, Pope, et al., 2016; Landaverde-González et al., 2017), suggesting that the resolution of these microsatellite-based studies only allowed the detection of very strong IBR effects. Our findings thus imply that fine-scale genetic structure and IBR may be more common in this group of bees than previously acknowledged and that studies employing thousands of genetic markers and large sample sizes are needed to identify or rule out weak, but significant IBR effects.

Our results reveal that thermal stability and forest cover (which were highly correlated) are key mediators of genetic connectivity in this stingless bee species, and support earlier findings stressing out the role of elevation as a bee dispersal barrier (Jackson et al., 2018; Jaffé, Pope, et al., 2016). Although thermal range is known to have a profound influence on insect physiology (Dixon et al., 2009), ours is the first study to report an effect on dispersal behaviors. Interestingly, temperature annual range was found to be the main abiotic predictor of bee richness and diversity in the eastern Neotropics, suggesting environmental variability may have led to higher speciation (Faria & Gonçalves, 2013). Our results imply that a similar mechanism may be operating at the species level, with increased thermal variability hindering gene flow, and thus facilitating local adaptation (Allendorf et al., 2013).

Even though thermal stability was the main factor explaining gene flow patterns in *M. subnitida*, temperature annual range and forest cover were highly correlated, so we could not disentangle the relative contribution of each. Our work is nevertheless the first to reveal a significant effect of forest cover on stingless bee gene flow, a long-standing expectation for this group of bees with restricted dispersal (Jaffé, Pope, et al., 2016). Reduced gene flow across deforested areas was only found in one other tropical organism so far (Monteiro et al., 2019), namely the army ant *Eciton burchelli* Westwood, 1842 (Pérez-Espona, McLeod, & Franks, 2012).



FIGURE 6 Spatial distribution of adaptive genetic variability in *Melipona subnitida*. Colors represent interpolated spatial principal components (sPCA) and suggest a latitudinal pattern associated with sPC1 (a) and an altitudinal pattern associated with sPC2 (b). Shaded areas represent elevations of at least 500 masl



Interestingly, these army ants exhibit striking life history similarities with stingless bees, as queens are permanently wingless and thus show a restricted dispersal (Jaffé, Moritz, & Kraus, 2009). Forested areas thus seem important dispersal corridors for this stingless bee, which could facilitate the migration toward higher elevations predicted under climate change (Supporting Information Figure S6).

Our environmental association tests can be considered robust to deviations from the underlying demographic structure, as candidate loci were intersected across different *k*-values. Additionally, LFMM were calibrated based on the distribution of adjusted *p*-values, so the incidence of false discovery rates was low (François et al., 2016). Interestingly, we found latitudinal and altitudinal gradients in the distribution of adaptive genetic variation. While the former gradient was also found in the honeybee *A. mellifera* (Chávez-Galarza et al., 2013; Henriques et al., 2018), and is probably related to climatic conditions, the altitudinal gradient suggests different adaptations to current highland and lowland areas. Our findings thus reveal the presence of

locally adapted bees, which should be preserved to maintain evolutionary potential (Hoffmann & Sgrò, 2011; Sgrò, Lowe, & Hoffmann, 2011). While lowland populations of *M. subnitida* are expected to shift to higher elevations by 2050 (Supporting Information Figure S6), current highland populations are at risk, since highlands will have no climate analogs in the future (Colwell, Brehm, Cardelús, Gilman, & Longino, 2008; Giannini et al., 2017). Conservation actions should thus prioritize the protection of current highland populations while improving the connectivity between highlands and lowlands, preserving or restoring foothill and mountain forests (Giannini et al., 2017).

Taken together, our findings shed important light on the life history of *M. subnitida* and highlight the role of regions with large thermal fluctuations, deforested areas, and mountain ranges as dispersal barriers. Moreover, our work unravels previously unknown patterns of local adaptation in these bees. This knowledge could help guide future conservation actions such as avoiding the transportation of colonies beyond 300 km, preserving highland and lowland populations Evolutionary Applications

separately, and conserving or restoring foothill and mountain forests. Considering the high biological and economic importance of this native pollinator, such conservation efforts will be easily offset if honey production and pollination services are maintained in the future.

DATA ARCHIVING STATEMENT

Genotype and sequence data have been deposited in FigShare: https://doi.org/10.6084/m9.figshare.7863416.v1

ACKNOWLEDGEMENTS

We thank Marcela de Matos Barbosa, Dirk Koedam, Jânio Ângelo Félix, Ulyses Madureira Maia, and Márcia Rêgo for help with the collection of specimens. R Jaffé was supported by a research grant from CNPq (478982/2013-5), JC Veiga by a CAPES/EMBRAPA PhD scholarship (15/2014), NS Pope by an NSF predoctoral fellowship, CS Carvalho by a CAPES postdoctoral fellowship (88887.156652/2017-00), and ECM Lanes by a CNPq postdoctoral fellowship (300714/2017-3). R Jaffé, BM Freitas, CF Martins, MC Arias, and VL Imperatriz-Fonseca received CNPg productivity research fellowship (301616/2017-5, 305126/2013-0, 302976/2015-9, 306932/2016-4, and 311531/2014-8, respectively). Funding was also provided by the Research Center on Biodiversity and Computing (BioComp) from the University of São Paulo, Empresa Brasileira de Pesquisa Agropecuária (02.11.01.029.00.00), and Instituto Tecnológico Vale.

CONFLICT OF INTEREST

None declared.

ORCID

Rodolfo Jaffé D https://orcid.org/0000-0002-2101-5282 Jamille C. Veiga D https://orcid.org/0000-0001-7554-2785 Éder C. M. Lanes 🕩 https://orcid.org/0000-0001-5326-0891 Carolina S. Carvalho D https://orcid.org/0000-0002-0063-2185 Ronnie Alves Dhttps://orcid.org/0000-0003-4139-0562 Maria C. Arias D https://orcid.org/0000-0003-1477-101X Airton T. Carvalho 🕩 https://orcid.org/0000-0003-3395-4444 Felipe A. L. Contrera D https://orcid.org/0000-0002-7078-5048 Tiago M. Francoy D https://orcid.org/0000-0002-2413-966X Breno M. Freitas D https://orcid.org/0000-0002-9932-2207 Tereza C. Giannini D https://orcid.org/0000-0001-9830-1204 Michael Hrncir Dhttps://orcid.org/0000-0003-4931-3924 Celso F. Martins D https://orcid.org/0000-0001-5953-3758 Guilherme Oliveira Dhttps://orcid.org/0000-0003-0054-3438 Vera L. Imperatriz-Fonseca ២ https://orcid. org/0000-0002-1079-2158

REFERENCES

- Ahrens, C. W., Rymer, P. D., Stow, A., Bragg, J., Dillon, S., Umbers, K. D. L., & Dudaniec, R. Y. (2018). The search for loci under selection: Trends, biases and progress. *Molecular Ecology*, 27(6), 1342–1356. https://doi. org/10.1111/mec.14549
- Alexander, D., Novembre, J., & Lange, K. (2009). Fast model-based estimation of ancestry in unrelated individuals. *Genome Research*, 19, 1655–1664. https://doi.org/10.1101/gr.094052.109
- Allendorf, F. W. (2017). Genetics and the conservation of natural populations: Allozymes to genomes. *Molecular Ecology*, *26*(2), 420–430. https://doi.org/10.1111/mec.13948
- Allendorf, F., Luikart, G., & Aitken, S. (2013). Conservation and the genetics of populations. Oxford, UK: Wiley-Blackwell.
- Alvarado-Serrano, D. F., Van Etten, M. L., Chang, S.-M., & Baucom, R. S. (2019). The relative contribution of natural landscapes and human-mediated factors on the connectivity of a noxious invasive weed. *Heredity*, 122(1), 29–40. https://doi.org/10.1038/s41437-018-0106-x
- Balkenhol, N., Cushman, S. A., Waits, L. P., & Storfer, A. (Eds.). (2016). In Landscape genetics. Hoboken, NJ: Chichester, UK. https://doi. org/10.1002/9781118525258
- Balkenhol, N., Dudaniec, R. Y., Krutovsky, K. V., Johnson, J. S., Cairns, D. M., Segelbacher, G., ... Selmoni, O. (2017). Landscape genomics: Understanding relationships between environmental heterogeneity and genomic characteristics of populations. In O. P.Rajora (Ed.), *Population genomics: Concepts, approaches and applications* (pp. 261– 322). Cham, Switzerland: Springer International Publishing. https:// doi.org/10.1007/13836_2017_2
- Barton, K. (2018). MuMIn: Multi-model inference. R package version 1.13.4.
- Benjamini, Y., & Hochberg, Y. (1995). Controlling the false discovery rate: A practical and powerful approach to multiple testing. Source Journal of the Royal Statistical Society. Series B (Methodological), 57(1), 289– 300. https://doi.org/10.2307/2346101
- Bonatti, V., Simões, Z. L. P., Franco, F. F., & Francoy, T. M. (2014). Evidence of at least two evolutionary lineages in *Melipona subnitida* (Apidae, Meliponini) suggested by mtDNA variability and geometric morphometrics of forewings. *Naturwissenschaften*, 101(1), 17–24. https://doi. org/10.1007/s00114-013-1123-5
- Brown, M. J. F. F., & Paxton, R. J. (2009). The conservation of bees: A global perspective. Apidologie, 40(3), 410-416. https://doi. org/10.1051/apido/2009019
- Bruno, C., Macchiavelli, R., & Balzarini, M. (2008). Non-parametric smoothing of multivariate genetic distances in the analysis of spatial population structure at fine scale. *Theoretical and Applied Genetics*, 117(3), 435–447. https://doi.org/10.1007/s00122-008-0788-1
- Chávez-Galarza, J., Henriques, D., Johnston, J. S., Azevedo, J. C., Patton, J. C., Muñoz, I., ... Pinto, M. A. (2013). Signatures of selection in the Iberian honey bee (*Apis mellifera iberiensis*) revealed by a genome scan analysis of single nucleotide polymorphisms. *Molecular Ecology*, 22(23), 5890–5907. https://doi.org/10.1111/mec.12537
- Clarke, R. T., Rothery, P., & Raybould, A. F. (2002). Confidence limits for regression relationships between distance matrices: Estimating gene flow with distance. *Journal of Agricultural*, *Biological*, and Environmental *Statistics*, 7(3), 361–372. https://doi.org/10.1198/108571102320
- Colwell, R. K., Brehm, G., Cardelús, C. L., Gilman, A. C., & Longino, J. T. (2008). Global warming, elevational range shifts, and lowland biotic attrition in the wet tropics. *Science*, 322, 258–261. https://doi. org/10.1126/science.1162547
- Cruz, D. O., Jorge, D. M. M., Pereira, J. O. P., Torres, D. C., Soares, C. E. A., Freitas, B. M., & Grangeiro, T. B. (2006). Intraspecific variation in the first internal transcribed spacer (ITS1) of the nuclear ribosomal DNA in *Melipona subnitida* (Hymenoptera, Apidae), an endemic stingless bee from northeastern Brazil. *Apidologie*, *37*, 376–386. https://doi. org/10.1051/apido

itionary Applicat

- Cushman, S. A., & Landguth, E. L. (2010). Spurious correlations and inference in landscape genetics. *Molecular Ecology*, 19(17), 3592–3602. https://doi.org/10.1111/j.1365-294X.2010.04656.x
- Danecek, P., Auton, A., Abecasis, G., Albers, C. A., Banks, E., DePristo, M. A., ... Durbin, R. (2011). Variant call format, binary variant call format and VCFtools. *Bioinformatics*, 27(15), 2156–2158. https://doi. org/10.1093/bioinformatics/btr330
- Davis, E., Murray, T., Fitzpatrick, U., Brown, M., & Paxton, R. (2010). Landscape effects on extremely fragmented populations of a rare solitary bee, *Colletes floralis*. *Molecular Ecology*, *19*, 4922–4935. https://doi.org/10.1111/j.1365-294X.2010.04868.x
- De Kort, H., Vandepitte, K., Bruun, H. H., Closset-Kopp, D., Honnay, O., & Mergeay, J. (2014). Landscape genomics and a common garden trial reveal adaptive differentiation to temperature across Europe in the tree species Alnus glutinosa. Molecular Ecology, 23(19), 4709–4721. https://doi.org/10.1111/mec.12813
- DiLeo, M. F., & Wagner, H. H. (2016). A landscape ecologist's agenda for landscape genetics. *Current Landscape Ecology Reports*, 1(3), 115– 126. https://doi.org/10.1007/s40823-016-0013-x
- Diniz-Filho, J. A. F., Nabout, J. C., Bini, L. M., Soares, T. N., de Campus Telles, M. P., de Marco, P., & Collevatti, R. G. (2009). Niche modelling and landscape genetics of Caryocar brasiliense ("Pequi" tree: Caryocaraceae) in Brazilian Cerrado: An integrative approach for evaluating central-peripheral population patterns. *Tree Genetics and Genomes*, 5, 617–627. https://doi.org/10.1007/ s11295-009-0214-0
- Dixon, A. F. G., Honěk, A., Keil, P., Kotela, M. A. A., Šizling, A. L., & Jarošík, V. (2009). Relationship between the minimum and maximum temperature thresholds for development in insects. *Functional Ecology*, 23(2), 257–264. https://doi.org/10.1111/j.1365-2435.2008.01489.x
- Do, C., Waples, R. S., Peel, D., Macbeth, G. M., Tillett, B. J., & Ovenden, J. R. (2014). NeEstimator v2: Re-implementation of software for the estimation of contemporary effective population size from genetic data. *Molecular Ecology Resources*, 14(1), 209–214. https://doi. org/10.1111/1755-0998.12157
- El-Niweiri, M. A. A., & Moritz, R. F. A. (2011). Mating in the rain? Climatic variance for polyandry in the honeybee (*Apis mellifera jemenitica*). Population Ecology, 53(3), 421-427. https://doi.org/10.1007/ s10144-011-0271-8
- Fahrig, L. (2013). Rethinking patch size and isolation effects: The habitat amount hypothesis. *Journal of Biogeography*, 40, 1649–1663. https:// doi.org/10.1111/jbi.12130
- Faleiro, F. V., Nemésio, A., & Loyola, R. (2018). Climate change likely to reduce orchid bee abundance even in climatic suitable sites. *Global Change Biology*, 24(6), 2272–2283. https://doi.org/10.1111/ gcb.14112
- Faria, L. R. R., & Gonçalves, R. B. (2013). Abiotic correlates of bee diversity and composition along eastern Neotropics. *Apidologie*, 44(5), 547–562. https://doi.org/10.1007/s13592-013-0205-x
- Ferreira, N. T., Blochtein, B., & Serrão, J. E. (2013). Seasonal production and spatial distribution of Melipona bicolor schencki (Apidae; Meliponini) castes in brood combs in southern Brazil. Apidologie, 44(2), 176-187. https://doi.org/10.1007/s13592-012-0169-2
- François, O., Martins, H., Caye, K., & Schoville, S. D. (2016). Controlling false discoveries in genome scans for selection. *Molecular Ecology*, 25, 454–469. https://doi.org/10.1111/mec.13513
- Frichot, E., & François, O. (2015). LEA: An R package for landscape and ecological association studies. *Methods in Ecology and Evolution*, 6(8), 925–929. https://doi.org/10.1111/2041-210X.12382
- Frichot, E., Mathieu, F., Trouillon, T., Bouchard, G., & François, O. (2014). Fast and efficient estimation of individual ancestry coefficients. *Genetics*, 196(4), 973–983. https://doi.org/10.1534/ genetics.113.160572
- Frichot, E., Schoville, S. D., Bouchard, G., & François, O. (2013). Testing for association between loci and environmental gradients using

latent factor mixed models. *Molecular Biology and Evolution*, 30(7), 1687–1699. https://doi.org/10.1093/molbev/mst063

- Giannini, T. C., Maia-Silva, C., Acosta, A. L., Jaffé, R., Carvalho, A. T., Martins, C. F., ... Imperatriz-Fonseca, V. L. (2017). Protecting a managed bee pollinator against climate change: Strategies for an area with extreme climatic conditions and socioeconomic vulnerability. *Apidologie*, 48(6), 784–794. https://doi.org/10.1007/ s13592-017-0523-5
- Hadley, A. S., & Betts, M. G. (2011). The effects of landscape fragmentation on pollination dynamics: Absence of evidence not evidence of absence. *Biological Reviews*, 87, 526–544. https://doi. org/10.1111/j.1469-185X.2011.00205.x
- Henriques, D., Wallberg, A., Chávez-Galarza, J., Johnston, J. S., Webster, M. T., & Pinto, M. A. (2018). Whole genome SNP-associated signatures of local adaptation in honeybees of the Iberian Peninsula. *Scientific Reports*, 8(1), 1–14. https://doi.org/10.1038/s41598-018-29469-5
- Hoffmann, A. A., & Sgrò, C. M. (2011). Climate change and evolutionary adaptation. *Nature*, 470(7335), 479–485. https://doi.org/10.1038/ nature09670
- Imperatriz-Fonseca, V., Koedam, D., & Hrncir, M. (2017). A abelha jandaíra: No passado, no presente e no futuro. EdUFERSA. Retrieved from https://www.researchgate.net/profile/Breno_Freitas/publication/ 321278052_Meliponineos_e_polinizacao_a_abelha_jandaira_ e_outros_meliponineos_na_polinizacao_agricola_no_ s e miarido/links/5a1868bfaca272df080a8388/ Meliponineos-e-polinizacao-aabelha-jandaira-e-
- Jackson, J. M., Pimsler, M. L., Oyen, K. J., Koch-Uhuad, J. B., Herndon, J. D., Strange, J. P., ... Lozier, J. D. (2018). Distance, elevation and environment as drivers of diversity and divergence in bumble bees across latitude and altitude. *Molecular Ecology*, 27(14), 2926–2942. https:// doi.org/10.1111/mec.14735
- Jaffé, R., Castilla, A., Pope, N., Imperatriz-Fonseca, V. L., Metzger, J. P., Arias, M. C., & Jha, S. (2016). Landscape genetics of a tropical rescue pollinator. *Conservation Genetics*, 17(2), 267–278. https://doi. org/10.1007/s10592-015-0779-0
- Jaffé, R., Moritz, R. F. A., & Kraus, F. B. (2009). Gene flow is maintained by polyandry and male dispersal in the army ant *Eciton burchellii*. *Population Ecology*, 51(2), 227–236. https://doi.org/10.1007/ s10144-008-0133-1
- Jaffé, R., Pioker-Hara, F. C., Dos Santos, C. F., Santiago, L. R., Alves, D. A., Kleinert, A. M. P., ... Imperatriz-Fonseca, V. L. (2014). Monogamy in large bee societies: A stingless paradox. *Naturwissenschaften*, 101(3), 261–264. https://doi.org/10.1007/s00114-014-1149-3
- Jaffé, R., Pope, N., Acosta, A. L., Alves, D. A., Arias, M. C., De la Rúa, P., ... Carvalheiro, L. G. (2016). Beekeeping practices and geographic distance, not land use, drive gene flow across tropical bees. *Molecular Ecology*, 25(21), 5345–5358. https://doi.org/10.1111/mec.13852
- Jaffé, R., Pope, N., Carvalho, A. T., Maia, U. M., Blochtein, B., de Carvalho, C. A. L., ... Imperatriz-Fonseca, V. L. (2015). Bees for development: Brazilian survey reveals how to optimize stingless beekeeping. *PLOS* ONE, 10(3), 1–21. https://doi.org/10.1371/journal.pone.0121157
- Jha, S. (2015). Contemporary human-altered landscapes and oceanic barriers reduce bumble bee gene flow. *Molecular Ecology*, 24(5), 993– 1006. https://doi.org/10.1111/mec.13090
- Jha, S., & Kremen, C. (2013). Urban land use limits regional bumble bee gene flow. *Molecular Ecology*, 22, 2483–2495. https://doi. org/10.1111/mec.12275
- Jombart, T. (2008). adegenet: A R package for the multivariate analysis of genetic markers. *Bioinformatics*, 24(11), 1403–1405. https://doi. org/10.1093/bioinformatics/btn129
- Kennedy, C. M., Lonsdorf, E., Neel, M. C., Williams, N. M., Ricketts, T. H., Winfree, R., ... Kremen, C. (2013). A global quantitative synthesis of local and landscape effects on wild bee pollinators in agroecosystems. *Ecology Letters*, 16(5), 584–599. https://doi.org/10.1111/ ele.12082

II FV Evolutionary Appli

- Kerr, J. T., Pindar, A., Galpern, P., Packer, L., Potts, S. G., Roberts, S. M., ... Pantoja, A. (2015). Climate change impacts on bumblebees converge across continents. *Science*, 349(6244), 177–180. https://doi. org/10.1126/science.aaa7031
- Koffler, S., Menezes, C., Menezes, P. R., Kleinert, A. D. M. P., Imperatriz-Fonseca, V. L., Pope, N., & Jaffé, R. (2015). Temporal variation in honey production by the stingless bee *Melipona subnitida* (Hymenoptera: Apidae): Long-term management reveals its potential as a commercial species in Northeastern Brazil. *Journal of Economic Entomology*, 108(3), 858–867. https://doi.org/10.1093/jee/tov055
- Landaverde-González, P., Enríquez, E., Ariza, M. A., Murray, T., Paxton, R. J., & Husemann, M. (2017). Fragmentation in the clouds? The population genetics of the native bee *Partamona bilineata* (Hymenoptera: Apidae: Meliponini) in the cloud forests of Guatemala. *Conservation Genetics*, 18(3), 631–643. https://doi.org/10.1007/ s10592-017-0950-x
- Le Conte, Y., & Navajas, M. (2008). Climate change: Impact on honey bee populations and diseases. *Revue Scientifique Et Technique*, 27, 499–510.
- Leroy, G., Carroll, E. L., Bruford, M. W., DeWoody, J. A., Strand, A., Waits, L., & Wang, J. (2018). Next-generation metrics for monitoring genetic erosion within populations of conservation concern. *Evolutionary Applications*, 11(7), 1066–1083. https://doi.org/10.1111/eva.12564
- Lozier, J. D. (2014). Revisiting comparisons of genetic diversity in stable and declining species: Assessing genome-wide polymorphism in North American bumble bees using RAD sequencing. *Molecular Ecology*, 23(4), 788–801. https://doi.org/10.1111/mec.12636
- Lozier, J., & Zayed, A. (2017). Bee conservation in the age of genomics. Conservation Genetics, 18(3), 713–729. https://doi.org/10.1007/ s10592-016-0893-7
- Maia-Silva, C., Hrncir, M., Silva, C. I., & Imperatriz-Fonseca, V. L. (2015). Survival strategies of stingless bees (*Melipona subnitida*) in an unpredictable environment, the Brazilian tropical dry forest. *Apidologie*, 46(5), 631–643. https://doi.org/10.1007/s13592-015-0354-1
- Marengo, J. A., Torres, R. R., & Alves, L. M. (2017). Drought in Northeast Brazil–past, present, and future. *Theoretical and Applied Climatology*, 129(3–4), 1189–1200. https://doi.org/10.1007/s00704-016-1840-8
- McCartney-Melstad, E., Vu, J. K., & Shaffer, H. B. (2018). Genomic data recover previously undetectable fragmentation effects in an endangered amphibian. *Molecular Ecology*, 27(22), 4430–4443. https://doi. org/10.1111/mec.14892
- McRae, B. H. (2006). Isolation by resistance. *Evolution*, 60(8), 1551. https://doi.org/10.1554/05-321.1
- Monteiro, W. P., Veiga, J. C., Silva, A. R., Carvalho, C. S., Lanes, É. C. M., Rico, Y., & Jaffé, R. (2019). Everything you always wanted to know about gene flow in tropical landscapes (but were afraid to ask). *PeerJ*, 7, e6446. https://doi.org/10.7717/peerj.6446
- Pedro, S. R. M. (2014). The stingless bee fauna in Brazil (Hymenoptera: Apidae). Sociobiology, 61, 348–354. https://doi.org/10.13102/sociobiology.v61i4.348-354
- Pérez-Espona, S., McLeod, J. E., & Franks, N. R. (2012). Landscape genetics of a top neotropical predator. *Molecular Ecology*, 21(24), 5969– 5985. https://doi.org/10.1111/mec.12088
- Pinheiro, J., Bates, D., DebRoy, S., Sarkar, D., & R Core Team (2018). nlme: Linear and nonlinear mixed effects models. R package version 3.1-137. Retrieved from https://cran.r-project.org/package=nlme
- Potts, S. G., Biesmeijer, J. C., Kremen, C., Neumann, P., Schweiger, O., & Kunin, W. E. (2010). Global pollinator declines: Trends, impacts and drivers. *Trends in Ecology and Evolution*, 25(6), 345–353. https://doi. org/10.1016/j.tree.2010.01.007
- Potts, S. G., Imperatriz-Fonseca, V. L., Ngo, H. T., Biesmeijer, J. C., Breeze, T. D., Dicks, L. V., ... Viana, B. F. (2016). *IPBES (2016): Summary for* policymakers of the assessment report of the Intergovernmental Science
 Policy platform on biodiversity and ecosystem services on pollinators, pollination and food production. Bonn, Germany: Secretariat of

the Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services.

- Pyke, G. H., Thomson, J. D., Inouye, D. W., & Miller, T. J. (2016). Effects of climate change on phenologies and distributions of bumble bees and the plants they visit. *Ecosphere*, 7(3), e01267. https://doi. org/10.1002/ecs2.1267
- Rellstab, C., Gugerli, F., Eckert, A. J., Hancock, A. M., & Holderegger, R. (2015). A practical guide to environmental association analysis in landscape genomics. *Molecular Ecology*, 24(17), 4348–4370. https:// doi.org/10.1111/mec.13322
- Romiguier, J., Gayral, P., Ballenghien, M., Bernard, A., Cahais, V., Chenuil, A., ... Galtier, N. (2014). Comparative population genomics in animals uncovers the determinants of genetic diversity. *Nature*, *515*(7526), 261–263. https://doi.org/10.1038/nature13685
- Roubik, D. W. (2006). Stingless bee nesting biology. Apidologie, 37(2), 124–143. https://doi.org/10.1051/apido
- Santos-Filho, P. S., Alves, D. A., Eterovic, A., Imperatriz-Fonseca, V. L., & Kleinert, A. M. P. (2006). Numerical investment in sex and caste by stingless bees (Apidae: Meliponini): A comparative analysis. *Apidologie*, 37, 207-221. https://doi.org/10.1051/apido:2006015
- Schlaepfer, D. R., Braschler, B., Rusterholz, H.-P., & Baur, B. (2018). Genetic effects of anthropogenic habitat fragmentation on remnant animal and plant populations: A meta-analysis. *Ecosphere*, 9(10), e02488. https://doi.org/10.1002/ecs2.2488
- Sgrò, C. M., Lowe, A. J., & Hoffmann, A. A. (2011). Building evolutionary resilience for conserving biodiversity under climate change. Evolutionary Applications, 4(2), 326–337. https://doi. org/10.1111/j.1752-4571.2010.00157.x
- Shirk, A. J., Landguth, E. L., & Cushman, S. A. (2017). A comparison of individual-based genetic distance metrics for landscape genetics. *Molecular Ecology Resources*, 17(6), 1308–1317. https://doi. org/10.1111/1755-0998.12684
- Silva, G. R., Souza, B. A., Pereira, F. M., Lopes, M. T. R., Valente, S. E., & Diniz, F. (2014). New molecular evidence for fragmentation between two distant populations of the threatened stingless bee Melipona subnitida Ducke (Hymenoptera, Apidae, Meliponini). *Journal of Hymenoptera Research*, *38*, 1–9. https://doi.org/10.3897/jhr.38.7302
- Silva, M. D., & Ramalho, M. (2016). The influence of habitat and species attributes on the density and nest spacing of a stingless bee (Meliponini) in the Atlantic Rainforest. *Sociobiology*, *63*(3), 991. https://doi.org/10.13102/sociobiology.v63i3.1037
- Storfer, A., Patton, A., & Fraik, A. K. (2018). Navigating the interface between landscape genetics and landscape genomics. *Frontiers in Genetics*, 9, 68. https://doi.org/10.3389/fgene.2018.00068
- Tajima, F. (1989). Statistical method for testing the neutral mutation. Genetics Society of America, 123, 585–595. Retrieved from http:// www.mathstat.concordia.ca/faculty/lpopovic/Tajima1989.pdf
- Tavares, M. G., Carvalho, C. R., Aparecida, F., & Soares, F. (2010). Genome size variation in *Melipona* species (Hymenoptera: Apidae) and subgrouping by their DNA content. *Apidologie*, 41, 636–642.
- Theodorou, P., Radzevičiūtė, R., Kahnt, B., Soro, A., Grosse, I., & Paxton, R. J. (2018). Genome-wide single nucleotide polymorphism scan suggests adaptation to urbanization in an important pollinator, the red-tailed bumblebee (Bombus lapidarius L.). Proceedings of the Royal Society B: Biological Sciences, 285(1877), 20172806. https://doi. org/10.1098/rspb.2017.2806
- Van Veen, J. W., & Sommeijer, M. J. (2000). Colony reproduction in Tetragonisca angustula (Apidae, Meliponini). Insectes Sociaux, 47(1), 70–75. https://doi.org/10.1007/s000400050011
- Viana, B. F., Boscolo, D., Mariano-Neto, E., Lopes, L. E., Lopes, A. V., Ferreira, P. A., ... Primo, L. M. (2012). How well do we understand landscape effects on pollinators and pollination services? *Journal of Pollination Ecology*, 7, 31–41.
- Vieira, R. M. S. P., Tomasella, J., Alvalá, R. C. S., Sestini, M. F., Affonso, A. G., Rodriguez, D. A., ... Santana, M. O. (2015). Identifying areas

WILE

susceptible to desertification in the Brazilian northeast. *Solid Earth*, *6*, 347–360. https://doi.org/10.5194/se-6-347-2015

- Vit, P., Pedro, S. R. M., & Roubik, D. (Eds.). (2013). In *Pot-honey*. New York, NY: Springer Science & Business Media. https://doi. org/10.1007/978-1-4614-4960-7
- Willmer, P. (2014). Climate change: Bees and orchids lose touch. Current Biology, 24(23), R1133–R1135. https://doi.org/10.1016/j. cub.2014.10.061
- Wratten, S., Gillespie, M., Decourtye, A., Mader, E., & Desneux, N. (2012). Pollinator habitat enhancement: Benefits to other ecosystem services. Agriculture, Ecosystems and Environment, 159, 112–122. https://doi.org/10.1016/j.agee.2012.06.020
- Yang, J., Benyamin, B., Mcevoy, B. P., Gordon, S., Henders, A. K., Dale, R., ... Visscher, P. M. (2010). Common SNPs explain a large proportion of heritability for human height. *Nature Genetics*, 42(7), 565–569. https://doi.org/10.1038/ng.608

SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section at the end of the article.

How to cite this article: Jaffé R, Veiga JC, Pope NS, et al. Landscape genomics to the rescue of a tropical bee threatened by habitat loss and climate change. *Evol Appl*. 2019;12:1164–1177. https://doi.org/10.1111/eva.12794